FEEDS & FEEDING

A TECHNICAL PUBLICATION OF THE ANIMAL NUTRITION ACADEMY
Volume 2 – Issue 34

Six trace minerals that influence eggshell quality

Ioannis Mavromichalis, MSc, PhD

This issue examines six trace minerals, their roles in eggshell formation, and the distinction between practical supplementation benchmarks and experimental interventions.

Eggshell quality remains an important economic characteristic of commercial egg production. Cracked and broken eggs represent direct losses, particularly as laying cycles extend and shells deteriorate with advancing hen age. Calcium receives most attention because the shell consists predominantly of calcium carbonate, but calcium alone cannot produce a strong shell. Formation also requires an organic matrix, sound membranes, and healthy reproductive tissues. Trace minerals contribute to these processes, although their mechanisms and the evidence supporting supplementation differ considerably. Evaluating the complete feeding program helps distinguish these possibilities before changing the premix, particularly when older hens produce larger eggs or feed intake falls.

Trace mineral sources include conventional inorganic compounds, organic forms, including amino acid complexes and chelates, and alternative inorganic sources. These differ in concentration, solubility, reactivity, stability, and biological availability; dietary interactions also affect utilization. Their relative advantages deserve a separate article. Here, supplementation benchmarks refer to elemental mineral added per kilogram of complete feed, excluding the contribution from ingredients. They summarize commercial laying-period recommendations rather than minimum requirements or guaranteed shell-quality responses.

Manganese builds the shell matrix

Manganese supports enzymes involved in synthesizing glycosaminoglycans and other components of the organic matrix that regulates calcium carbonate deposition. Deficiency can disrupt shell ultrastructure and reduce strength even when calcium intake is adequate. Most authorities converge on 100 mg supplemental Mn/kg for laying hens. Supplementation has improved shell characteristics in some experiments, but responses depend on basal supply, source, and flock conditions. Even in aged hens, improved production does not necessarily coincide with improved breaking strength. Correcting inadequate manganese supply is therefore different from increasing an already adequate specification.

Zinc supports shell calcification

Zinc is a component of carbonic anhydrase, which catalyzes the conversion of carbon dioxide and water into bicarbonate. This provides substrate for the carbonate incorporated into the shell. Zinc also supports proteins involved in mineralization and crystal organization. Recommendations span 60–80 mg supplemental Zn/kg. Inadequate supply can compromise calcification despite satisfactory calcium intake. However, supplementation responses depend on existing supply, availability, and hen age. A nutritional role does not establish that progressively increasing zinc will continue to improve shells.

Copper supports shell membranes
Copper contributes to the structural integrity of shell membranes, which provide the framework for shell mineralization. It is required by lysyl oxidase, an enzyme involved in cross-linking structural proteins. Deficiency can therefore impair normal shell development. Most authorities about 10 mg supplemental Cu/kg for laying hens. Trials combining copper with manganese and zinc have reported improvements but cannot attribute every response to copper alone. Excessive supplementation can disturb mineral balance.

Selenium supports cell protection
Selenium contributes mainly through selenoproteins, including glutathione peroxidases, that protect cells against oxidative damage. This supports reproductive tissue function without making selenium a major structural constituent of the shell. Breeder supplementation benchmarks span 0.20–0.30 mg Se/kg. Some experiments in aged, selenium-depleted hens have improved shell characteristics, but other trials found no improvement in shell thickness or strength. Results from deficient birds should not be assumed to apply to adequately supplemented commercial flocks. Selenium supplementation requires accurate mixing and attention to the permitted source, supplemental allowance, and total dietary concentration under applicable regulations.

Iron supports general metabolism
Iron supports oxygen transport, cellular respiration, and enzyme systems in the tissues responsible for egg production. Its direct contribution to shell structure is less clearly established than that of manganese, zinc, or copper. Selected breeder guides specify 25–40 mg supplemental Fe/kg. Ingredients often contribute substantial iron, but measured concentration alone does not establish availability. Additional iron should therefore be assessed against ingredient supply and mineral interactions.

Boron remains experimental
Boron has been investigated for effects on calcium metabolism, bone mineralization, and eggshell quality, but has no established commercial-layer requirement or breeder supplementation range. One White Leghorn experiment using 40 mg supplemental B/kg reported improved calcium retention and shell thickness, with fewer cracked eggs than the calcium-inadequate treatment. This is a research dose, not a commercial recommendation. Other experiments have produced variable responses, and high supplementation can depress intake and production. Boron therefore remains an experimental intervention requiring further evidence, rather than a routine replacement for established mineral nutrition.

In Brief, strong shells require adequate calcium and balanced trace-mineral nutrition. Manganese, zinc, and copper have established roles in shell formation; selenium and iron provide broader physiological support, while boron remains experimental. Breeder benchmarks guide supplementation, but increasing inclusion beyond an adequate supply does not guarantee better shells.

DISCLAIMER *Feeds & Feeding* is published by Animal Nutrition Academy for educational and reference purposes. Although reasonable care is taken to ensure accuracy, the author(s) and Animal Nutrition Academy accept no liability, to the extent permitted by law, for loss or damage arising from reliance on or application of this material. Readers remain responsible for its interpretation and use.

COPYRIGHT NOTICE © Animal Nutrition Academy. All rights reserved. No part of this publication may be reproduced, stored, or transmitted in any form without prior written permission. www.animalnutritionacademy.com.